

**PRELIMINARY PHYTOCHEMICAL ANALYSIS OF THE FERN
CHEILOSORIAMYSURENSIS(WALL. EX HOOK.)
CHING & SHING AND ITS ANTIOXIDANT ACTIVITY**

R. Rakkimuthu*, D. Sowmiya and P. Ananthi

*PG and Research Department of Botany
Nallamuthu Gounder Mahalingam College, Pollachi-642001
* Corresponding Author: biorakki@gmail.com*



Abstract

Cheilosoriamysurensis is a medicinal fern used to treat a variety of ailments, with few researches done on its phytochemical makeup and potential bioactivity. The main aim of the study was to support that traditional herbal medicine with active phytochemicals would be a good substitute for allopathic medicine. The methanol extract was tested for preliminary phytochemical analysis and its antioxidant activity. *Cheilosoriamysurensis* exhibited significant reducing power and total antioxidant activity, and its absorbance increased with an increase in concentration of sample. Overall, the findings suggest that this fern could be useful for medical purposes as a source of phytochemicals.

Keywords: *Cheilosoriamysurensis*; antioxidant; medicinal fern; Reducing power: Total antioxidant assay:

Introduction

focuses on locally available plant species and plant-based products, as well as ancient knowledge, traditional medicine can be found all over the world (AwasandDemissew, 2009). Human society faces numerous obstacles, and coping with health problems is a serious societal concern, mainly in growing international locations with restricted resources (Mangambu, 2013). The development in biology and medicine, many people in growing countries do not have sufficient fitness care (Singh and Singh, 2012). The world population, especially the rural population in developing countries are still largely dependent on herbal treatments (Islam, 2014).

A number of diseases have become increasingly important in recent times, of which infectious diseases caused by bacteria and cancer are two areas that need immediate attention. Plant sources with diverse natural bioactive ingredients play an important role in curing dreaded diseases (Atanasov, *et al.*, 2015). Many chemical compounds show impressive in vitro activity against pathogenic microorganisms resistant to recent allergenic drugs. The identification and characterization of these lead molecules in terms of their biological activity on the spectrum, potency, toxicity and safety is most needed.

Pharmacological research and traditional medicine have brought many drugs into the international pharmacopoeia. Pteridophytes are immune to many microbial diseases, it is possible that this is one of the reasons for their evolutionary success and the fact that they have been around for over 350 million years. Screening plant extracts for antibacterial, antioxidant, and anti-cancer effects could be beneficial to both humans and plants, given the enormous diversity of Indian medicinal plants and pteridophytes. In recent years, antimicrobial resistance has become a global issue. Because medicinal ferns are biodegradable, harmless and have fewer side effects, it is necessary to screen them for novel bioactive chemicals (Prusti, 2008).

Recently, biological research has concentrated on the medicinal, pharmaceutical, and phytochemical aspects of pteridophytes, which have significant value for medical and industrial uses. Many pteridophyte species are still being studied for possible uses and for the extraction of new active components. Hence, in this work is carried out the preliminary phytochemical analysis and its antioxidant activity of the fern, *Cheilosoria maysurensis*, found in the Western Ghats, Tamilnadu.

Materials and Methods

Plant material

Cheilosoria maysurensis (wall. ex hook.) ching & K. Hsing were collected from the Gopala Swamy hills, Western Ghats of Tamilnadu, South India. The fern species was identified and authenticated by BSI, Southern circle, Coimbatore, Tamilnadu. A voucher specimen was reposed as herbarium at the Department of Botany, Nallamuthu Gounder Mahalingam College, Pollachi.

Sample preparation

The methanolic extract of the fern leaves (Dried) were prepared according to the method described by Harbone, 1998. The sample was extracted using a soxhlet apparatus. The collected extract was concentrated under reduced pressure at 40°C. The extract was dried and stored at 4°C in storage vials for experimental use.

Qualitative phytochemical Analysis

Test for alkaloids

About 2 g of plant materials were crushed then added 1 mL of ammonia. Furthermore, 10 mL of chloroform was added, then crushed and filtered. The filtrate was added 10 mL of sulfuric acid 2N, shaken vigorously, left for a minute until the sulfuric acid solution and chloroform separated. The sulfuric acid layer is taken into a test tubes and tested by Meyer reagents to determine the presence of alkaloids. The addition of Meyer reagent established white precipitate indicate the presence of alkaloids (Miles, 1985).

Test for Terpenoid, Steroid, and Saponin:

The methanol extract was concentrated and partitioned with hexane. The soluble extract in hexane was tested with the Liberman-Bourchard reagent. The blue or green color exhibits the presence of steroids and red color for terpenoids. The insoluble residue in hexane is added water and shaken vigorously. The presence of the stable foam for 30 min indicates the existence of saponins, if positive for saponins, the solution was hydrolyzed with HCl and tested with the Liberman-Bourchard reagent. The green or blue color indicates the presence of steroid saponins and the purple or red color shows the existence of terpenoid saponins (Lajis, 1994).

Test for Flavonoid:

The methanol extract was concentrated and partitioned with hexane. The residue was extracted with 10 mL of 80% ethanol, subsequently added 0.5 mg of magnesium and HCl 0.5 M. The pink or purple color shows the presence of flavonoids (Halimatussakdiah and Amna, 2016)

Test for Phenol:

Methanol extract tested by Ferric Chloride. Add 3 – 4 drops of FeCl₃ solution into extract, the formation of bluish black color exhibits the phenol compound (Tiwari *et al.*, 2011).

Test for Tannin:

About 0.5 g of methanol extract was boiled in 10 ml of water in the test tube and then filtered. Add a few drops of FeCl₃ 0.1%. Forming of a brownish green or bluish black color indicates tannins (Ayoola *et al.*, 2008)

Test for Glycosides (Keller-Killani Test):

Glacial acetic acid was added into 2 ml. extract and one drop 5% FeCl₃ and conc. H₂SO₄. Reddish brown color appears at the junction of the two liquid layers and the upper layer of bluish green indicates the presence of glycosides (Kokate *et al.*, 2001).

Antioxidant activity

Sample preparation

The methanol leaf extract of *Cheilosoria myrsurensis* 1.0 mg was dissolved in 1 ml of DMSO. From the above 125, 250, 500 and 1000 µg/ml concentration of sample was prepared and used for different assays. Ascorbic acid was used as standard for comparing the activities.

Reducing power assay

The reducing power of methanolic leaf extracts of *C. myrsurensis* was determined according to the method previously described by Oyaizu (1986). Different concentrations of methanol extracts of *C.*

mysurensis was mixed with phosphate buffer (2.5ml, 0.2M, pH 6.6) and potassium ferricyanide [$K_3Fe(CN)_6$] (2.5 ml, 1%) separately. The mixtures were incubated at 50°C for 20 min. Then, 2.5 ml of trichloroacetic acid (10%) was added to the mixtures and centrifuged at 3000 rpm for 10 min. The upper layer of the solution (2.5 ml) was mixed with distilled water (2.5ml) and $FeCl_3$ (0.5 ml, 0.1%) and the absorbance was measured at 700 nm. Ascorbic acid was used as the standard. Phosphate buffer (pH 6.6) was used as blank.

Total antioxidant capacity by phosphomolybdenum assay

The spectrophotometric measurement of Total antioxidant capacity (TAC) is based on the reduction of Mo (VI) to Mo (V) by antioxidant compound and the formation of green phosphate / Mo (v) complex at acidic pH (Prieto *et al.*, 1999). Different concentrations (125- 1000 μ g/ml) of methanolic leaf extracts were combined with 3 ml of reagent solution (0.6 M sulfuric acid, 28 mM sodium phosphate and 4mM ammonium molybdate). The tubes containing the reaction solution was incubated at 95°C for 90 min. Then the absorbance of the solution was measured at 695 nm. Methanol (0.3 ml) in the place of extract was used as the blank. Ascorbic acid was used as a reference standard.

Results and Discussion

Phytochemical analysis:

The preliminary phytochemical analysis results of *Cheilosoria mysurensis* methanol extract was recorded (Table-1). *C. mysurensis* contains Alkaloids, steroids, saponins, phenols tannins and coumarins.

Table-1 Phytochemical analysis of methanol extract of *Cheilosoria mysurensis*

Secondary Metabolites	Methanol Extract of <i>Cheilosoria mysurensis</i>
Alkaloid	+
Terpenoid	-
Steroid	+
Saponin	+
Flavonoid	-
Phenol	+
Tannin	+
Glycosides	+

Pradnya *et al.* (2015) reported phytochemical analysis of four *Cheilanthes* species from northern Western Ghats of India. The methanolic extract of these ferns contains steroids, triterpenoids, reducing sugar, sugars, alkaloids, phenolic compounds, flavonoids, catechins and saponins. In our present investigation methanol extract of *Cheilosoria mysurensis* expressed the positive results for various phytochemical results. Pandey *et al.* (2014) investigated phytochemical analysis of methanolic extract

of *Adiantum* and *Pteris* leaf and stem contains tannins, saponins, flavonoids, and terpenoids steroid terpenoid. Smiliar results were obtained in the present study.

Damaynatiet *al.* (2019) reported secondary metabolites like alkaloids, flavonoids, carbohydrates, glycosides, phenol, saponin, steroid, tannins were present in the methanol extract of *Aspleniumindicum*, *Lepisorusnudum* and *Microsorummembranecium* . In the present study methanol extract of *Cheilosoriamsurensis* expressed the positive results for various phytochemical results.Bharti (2018) reported that methanolic extract of *Ampelopterisprolifera* and *Lygodiumflexosum* shows phytochemical compounds of carbohydrates, protein, aminoacids, alkaloids, glycosides, saponins,flavonoids, phenolics and tannins. In the present study methanol extract of *Cheilosoriamsurensis* expressed the positive results for various phytochemical results.

Herinet *al.* (2013) reported the methanolic extract of *Pterisargyreae*, *P.confuse*, *P.vittata* , *P. biaurita*and *P. multiaurita* shows phytochemical compounds of steroids, alkaloids, saponins , flavonoids, triterpenoids, phenolic compounds and tannins. In present study methanol extract of *Cheilosoriamsurensis* results were directly coincided with the above results.

Antioxidant activity

Reducing power assay

The reducing power of *C. mysurensis* methanol extract was shown in **Table 2**. The reducing power of methanol extract of absorbance (0.10 ± 0.01 to 0.03 ± 0.006) increased with concentration of methanol extracts of $12.5 \mu\text{g/ml}$ to $100 \mu\text{g/ml}$. The *C. mysurensis* extract showed potent ferric reducing power at $100 \mu\text{g/ml}$ concentration. Ascorbic acid had increase in absorbance value indicate increase in Reducing power capacity.

How yeelai *et al.* (2011) evaluated that antioxidant activities of the methanolic extracts of selected ferns in Malaysia. In this FRP method measured the ability of an antioxidant to donate electron to Fe (III) resulting in the reducing of Fe^{3+} / ferricynaide complex to Fe^{2+} complex, which would be monitored at 700 nm. Results we expressed as Gallic acid equivalent (GAE)/100 g leaves. The higher the FRP value, the greater is the reducing power of tested compound, thus greater the antioxidant activity. In the present study, the results indicate that the reducing ability of the extracts increased with the concentration. The result of *C. mysurensis* was significant. The reducing power of *C. mysurensis* extracts in the descending order. The reducing capacity of ascorbic acid was found to be higher than the extracts at each concentration points.

Gupta *et al.* (2014) reported the In vitro antioxidant activity of methanolic extracts of some ferns from Mawsynram of Meghalaya, India. The reducing ability of *Aleuritopterisflava* extracts was determined with ascorbic acid equivalent. Higher ascorbic acid equivalent value indicates higher reducing capacity

of sample, thus greater antioxidant potential. In the present study, the results indicate that the reducing ability of the extracts of *C. mysurensis* was increased with the ascorbic acid. Pandey *et al.* (2014) reported that antioxidant potential of methanolic extract of Adiantum and Pteris ferns. Analysis of leaf of Adiantum and Pteris ferns has a more 33 reducing power compared to their stem. There is a significant increase in absorbance of the reaction mixture indicates reducing power. In the present study, the result indicate that the reducing ability of the extracts of *C. mysurensis* was increased with the absorbance.

Table 2: Reducing power assay of *C. mysurensis*

Concentration	12.5 $\mu\text{g/ml}$	25 $\mu\text{g/ml}$	50 $\mu\text{g/ml}$	100 $\mu\text{g/ml}$
Methanol extract (Absorbance)	0.03 \pm 0.006	0.07 \pm 0.03	0.09 \pm 0.01	0.10 \pm 0.01
Concentration	12.5 $\mu\text{g/ml}$	25 $\mu\text{g/ml}$	50 $\mu\text{g/ml}$	100 $\mu\text{g/ml}$
Ascorbic acid (Absorbance)	0.101	0.109	0.374	0.762

Total antioxidant capacity assay

Increase in absorbance indicates increase in total antioxidant capacity. The methanolic extract of *C. mysurensis* exhibited significant activity and its absorbance increased with increase in concentration and the values were in comparison with ascorbic acid. The methanol extract exhibited significant absorbance of 2.1 ± 0.14 for $100\mu\text{g/ml}$. Ascorbic acid had an absorbance 0.74 at $100 \mu\text{g/ml}$. Jose *et al.* (2017) studied the antioxidant effects of hexane, ethyl acetate, and methanolic extract of *Pyrrosia heterophylla*. The assay is based on reduction of phosphate – molybdenum (IV) to phosphate molybdenum (V) by various concentrations (50,100,200,400 and $800\mu\text{g/ml}$). The result of *C. mysurensis* also show similar activity.

Saleha *et al.* (2014) reported the total antioxidant activity of fern *Diplazium esculentum*. The total antioxidant potentials of DECH and DEM showed higher antioxidant activity compared to standard ascorbic acid. In the present study, methanolic extract of *C. mysurensis* also show the same result. Selvaraj *et al.* (2015) revealed the antioxidant effects of methanolic extract of *Azolla microphylla*. From this analysis, it was evident that antioxidant increases with increase in concentration. The result of *C. mysurensis* exhibited significant activity and also antioxidant increases with increase in concentration. The total antioxidant capacity of *C. mysurensis* was shown in Table 3.

Table 3: Total antioxidant capacity by phosphomolybednum of *C. mysurensis*

Concentration	12.5 μ g/ml	25 μ g/ml	50 μ g/ml	100 μ g/ml
Methanol extract (Absorbance)	0.4 \pm 0.04	0.7 \pm 0.05	1.2 \pm 0.02	2.1 \pm 0.14
Concentration	12.5 μ g/ml	25 μ g/ml	50 μ g/ml	100 μ g/ml
Ascorbic acid (Absorbance)	0.068	0.158	0.313	0.700

Conclusion

The chemical composition, antioxidant, and cytotoxicity activities of the methanol extract of *C. mysurensis* from the Western Ghats of South India are reported for the first time. According to the findings of this study methanol extract of the fern contains contain Alkaloids, steroids, saponins, phenols tannins and coumarins. Furthermore, the findings demonstrated that the *C. mysurensis* methanol extract contains powerful antioxidant activities, as well as being a promising source for developing future antioxidant and anticancer medications.

Reference

1. Akter, Saleha& Hossain, Md&Ara, Ismot& Akhtar, Parvez. (2014). Investigation of in vitro antioxidant, antimicrobial and cytotoxic activity of *Diplaziumesculentum* (Retz). SW.. International Journal of Advances in Pharmacy, Biology and Chemistry. 3.
2. Atanasov, A. G. *et al.*,(2015). Discovery and resupply of pharmacologically active plant-derived natural products: a review. Biotechnol. Adv. 33, 1582–1614.
3. Awadhesh Kumar, Shrivastavaa, Bhawana, Pandey B and DeeptiChauhan, (2014).Phytochemical analysis of ferns Adiantum and Pteris ferns and its role as antioxidantIndianJ.Sci.Res. (4) Page no. 31-38.
4. Awas T, and Demissew S (2009). Ethno botanical study of medicinal plants in Kafficho people, south western Ethiopia. In Proceedings of the 16th International Conference of Ethiopian Studies. (3): 711-726.
5. Ayoola, G. A., Coker, H. A. B., Adesegun, S. A., Adepoju-Bello, A. A., Obaweya, K., Ezennia, E. C., and Atangbayila, T. O. (2008). Phytochemical Screening and Antioxidant Activities of Some Selected Medicinal Plants Used for Malaria Therapy in Southwestern Nigeria. Tropical Journal of Pharmaceutical Research, 7(3), 1019- 1024.
6. Bharti, (2018).Studied the phytochemical analysis and screening for active compounds in some ferns of Ranchi and Latehar districts, International journal of academic research and development (3) page no. 33-41.
7. DamaynatiJadhav, Mandaghatage and Vanitakarande, (2018).Phytochemicals studies on three epiphytic ferns from Mahabaleshwar and Panchgani hills. RJLBCS (5) page no.680-686.

8. Gupta SK, Ghosal M, Biswas R, Saha BK, Das AP and Mandal P. (2014). Evaluation of in vitro antioxidant activity of methanolic extracts of some ferns from Mawsynram of Meghalaya, India. *Int J Curr Sci.*, 12; E 87-97.
9. Halimatussakdiah and Amna, U. (2016). IsolasiSenyawa Alkaloid IndoldariEkstrakAkar*Kopsiasingapurensis*Ridl. (Apocynaceae). *JurnalIlmiah JURUTERA*, (3)1, 032-037.
10. Harborne, J.B. (1998). *Textbook of Phytochemical Methods. A Guide to Modern Techniques of Plant Analysis.* 5th Edition, Chapman and Hall Ltd, London, 21-72.
11. Herinsheebagracelin, D., John de britto,A and P. Benjamin Jeyarathnakumar, (2013).Qualitative and quantitative analysis of phytochemicals in five *pteriss*species, *International journal of pharmacy and pharmaceutical sciences*,(5) page no. 105-107.
12. How Yee Lai and YauYan Lim, (2011). Evaluation of antioxidant activities of the methanolic extracts of selected ferns in Malaysia, *International journal of environmental science and development* (2). Page no. 442-447
13. Islam MK, Saha S, Mahmud I, Mohamad K, Awang K, Uddin SJ, Rahman MM, Shilpi JA (2014). An ethno botanical study of medicinal plants used by tribal and native people of Madhupur forest area, Bangladesh. *Journal of Ethno pharmacology*; 151(2):921-930.
14. Jose kumar,V.S., Leenamma joseph and Meera George, (2017). Antioxidant, Antilipidperoxidative and antimicrobial properties of the epiphytic fern, *Pyrrosiaheterophylla* (L.) M.G.Price, *International journal of pharmacy and pharmaceutical sciences* (9), page no. 87-93.
15. Kokate C K, Purohit A P and Gokhale SB. (2001) Carbohydrate and derived Products, drugs containing glycosides, drugs containing tannins, lipids and protein alkaloids. *Text book of Pharmacognosy*, 7, edition: 133 -166, 167- 254, 255-2 69, 272- 310, 428-52 3.
16. Kunjiappanselvaraj, Bhattacharjeechiranjib and Chowdhury ranjana, (2015). Hepaprotective and antioxidant effect of *Azollamicrophylla* based gold nanoparticles against acetaminophen induced toxicity in a fresh water common crap fish (*Cyprinuscarpo* L.), *Nanomedicine journal* (2) page no. 88 – 110.
17. Lajis, S. M. K. (1994). The Phytochemical Survey, Proceeding of Workshop. Department Of Chemistry University Of Malaysia, Malaysia. LitbangKehutanan, Jakarta
18. Mangambu M (2013). Taxonomy, Biogeography and Ecology of the Pteridophytes of the mountain forest ecosystem of the KahuziBiega National Park in eastern DR Congo. Thesis, University of Antwerp, Belgium. p. 463.
19. Mathur A, and Joshi H (2012). Traditional remedies in Tarai region of Kumaun, Uttarakhand. *Indian Journal of Traditional Knowledge*; 11(4): 652-657.
20. Miles, D. H. (1985). *A Guide to Biologically Active Plants Constituent.* Departement of Chemistry, University of Flourida, USA

21. Oyaizu, M. (1986) Studies on Products of Browning Reactions: Antioxidative Activities of Product of Browning Reaction Prepared from Glucosamine. *Japan Journal of Nutrition*, 44, 307-315.
22. Pradnya N. GhorpadeS, B. Thakar, M.M. DongareANDManisha V. Kale. 2015, phytochemical analysis of four *cheilanthes* species from northern western Ghats of India, RJLBPCS(1) page no. 93-100.
23. Prieto P, Pineda M, Aguilar M. (1999). Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: Specific application to the determination of vitamin E. *Anal Biochem*. ;269:337–341.
24. Prusti A, Mishra SR, Sahoo S and Mishra SK (2008). Antibacterial Activity of some Indian Medicinal Plants. *Ethnobot Leaflets*, 12: 227-230.
25. Singh, Sand Singh R (2012). Ethno medicinal use of Pteridophytes in reproductive health of tribal women of Pachmarhi Biosphere Reserve, Madhya Pradesh, India. *International Journal of Pharmaceutical Sciences and Research*, 3(12): 4780.
26. Tiwari, P., Kumar. B., Kaur, M., Kaur, G., and Kaur, H. (2011). Phytochemical screening and Extraction: A Review. *InternationalePharmaceuticaSciencia* 1(1), 98-106. 25.